

4726 Further Pure Mathematics 2

1	(i)	Get $f'(x) = \pm \sin x / (1 + \cos x)$	M1	Reasonable attempt at chain at any stage
		Get $f''(x)$ using quotient/product rule	M1	Reasonable attempt at quotient/product
		Get $f(0) = \ln 2, f'(0) = 0, f''(0) = -1/2$	B1	Any one correct from correct working
			A1	All three correct from correct working
	(ii)	Attempt to use Maclaurin correctly	M1	Using their values in $af(0) + bf'(0)x + cf''(0)x^2$; may be implied
		Get $\ln 2 - 1/4 x^2$	A1✓	From their values; must be quadratic
2	(i)	Clearly verify in $y = \cos^{-1} x$	B1	i.e. $x = 1/2\sqrt{3}, y = \cos^{-1}(1/2\sqrt{3}) = 1/6\pi$, or similar
		Clearly verify in $y = 1/2 \sin^{-1} x$	B1	Or solve $\cos y = \sin 2y$
			SR	Allow one B1 if not sufficiently clear detail
	(ii)	Write down at least one correct diff' al	M1	Or reasonable attempt to derive; allow \pm
		Get gradient of -2	A1	cao
		Get gradient of 1	A1	cao
3	(i)	Get y - values of 3 and $\sqrt{28}$	B1	
		Show/explain areas of two rectangles equal y - value x 1 , and relate to A	B1	Diagram may be used
	(ii)	Show $A > 0.2(\sqrt{1+2^3} + \sqrt{1+2.2^3} + \dots$ $\dots \sqrt{1+2.83})$ $= 3.87(28)$	M1	Clear areas attempted below curve (5 values)
		Show $A < 0.2(\sqrt{1+2.2^3} + \sqrt{1+2.4^3} + \dots$ $\dots + \sqrt{1+3^3})$ $= 4.33(11) < 4.34$	A1	To min. of 3 s.f.
			M1	Clear areas attempted above curve (5 values)
			A1	To min. of 3 s.f.
4	(i)	Correct formula with correct r	M1	May be implied
		Expand r^2 as $A + B \sec \theta + C \sec^2 \theta$	M1	Allow $B = 0$
		Get $C \tan \theta$	B1	
		Use correct limits in their answer	M1	Must be 3 terms
		Limits to $1/12\pi + 2 \ln(\sqrt{3}) + 2\sqrt{3}/3$	A1	AEEF; simplified
	(ii)	Use $x = r \cos \theta$ and $r^2 = x^2 + y^2$	B1	Or derive polar form from given equation
		Eliminate r and θ	M1	Use their definitions
		Get $(x - 2)\sqrt{x^2 + y^2} = x$	A1	A.G.

4726

Mark Scheme

January 2008

5	(i)	Attempt use of product rule	M1	
		Clearly get $x = 1$	A1	Allow substitution of $x = 1$
	(ii)	Explain use of tangent for next approx. Tangents at successive approx. give $x > 1$	B1 B1	Not use of G.C. to show divergence Relate to crossing x -axis; allow diagram
6	(iii)	Attempt correct use of N-R with their derivative	M1	
		Get $x_2 = -1$	A1	✓
		Get $-0.6839, -0.5775, (-0.5672\dots)$	A1	To 3 d.p. minimum
		Continue until correct to 3 d.p.	M1	May be implied
		Get -0.567	A1	cao
6	(i)	Attempt division/equate coeff.	M1	To lead to some $ax + b$ (allow $b = 0$ here)
		Get $a = 2, b = -9$	A1	
		Derive/quote $x = 1$	B1	Must be equations
	(ii)	Write as quadratic in x	M1	$(2x^2 - x(11 + y) + (y - 6) = 0)$
		Use $b^2 \geq 4ac$ (for real x)	M1	Allow $<, >$
		Get $y^2 + 14y + 169 \geq 0$	A1	
		Attempt to justify positive/negative	M1	Complete the square/sketch
		Get $(y + 7)^2 + 120 \geq 0$ – true for all y	A1	
			SC	Attempt diff; quot./prod. rule M1
				Attempt to solve $dy/dx = 0$ M1
7	(i)	Get $x(1 + x^2)^{-n} - \int x \cdot (-n(1 + x^2)^{-n-1} \cdot 2x) dx$	M1	Reasonable attempt at parts
		Accurate use of parts	A1	
		Clearly get A.G.	B1	Include use of limits seen
(ii)	Express x^2 as $(1 + x^2) - 1$			
	Get $\frac{x^2}{(1 + x^2)^{n+1}} = \frac{1}{(1 + x^2)^n} - \frac{1}{(1 + x^2)^{n+1}}$	B1	Justified	
	Show $I_n = 2^{-n} + 2n(I_n - I_{n+1})$	M1	Clear attempt to use their first line above	
	Tidy to A.G.	A1		
(iii)	See $2I_2 = 2^{-1} + I_1$	B1		
	Work out $I_1 = \frac{1}{4}\pi$	M1	Quote/derive $\tan^{-1}x$	
	Get $I_2 = \frac{1}{4} + \frac{1}{8}\pi$	A1		

4726

Mark Scheme

January 2008

8	(i)	Use correct exponential for $\sinh x$	B1	
		Attempt to expand cube of this	M1	Must be 4 terms
		Correct cubic	A1	
		Clearly replace in terms of \sinh	B1	(Allow $\text{RHS} \rightarrow \text{LHS}$ or $\text{RHS} = \text{LHS}$ separately)
	(ii)	Replace and factorise	M1	Or state $\sinh x \neq 0$
		Attempt to solve for $\sinh^2 x$	M1	(= $\frac{1}{4}(k-3)$) or for k and use $\sinh^2 x > 0$
		Get $k > 3$	A1	Not \geq
	(iii)	Get $x = \sinh^{-1} c$	M1	($c = \pm \frac{1}{2}$); allow $\sinh x = c$
		Replace in \ln equivalent	A1√	As $\ln(\frac{1}{2} + \sqrt{\frac{5}{4}})$; their x
		Repeat for negative root	A1√	May be given as neg. of first answer (no need for $x=0$ implied)
			SR	Use of exponential definitions
				Express as cubic in $e^{2x} = u$ M1
		Factorise to $(u-1)(u^2-3u+1)=0$ A1		
		Solve for $x=0, \frac{1}{2}\ln(\frac{3}{2} \pm \sqrt{\frac{5}{2}})$ A1		
9	(i)	Get $\sinh y \frac{dy}{dx} = 1$	M1	Or equivalent; allow \pm Allow use of \ln equivalent with Chain Rule
		Replace $\sinh y = \sqrt{\cosh^2 y - 1}$	A1	
		Justify positive grad. to A.G.	B1	e.g. sketch
	(ii)	Get $k \cosh^{-1} 2x$	M1	No need for c
		Get $k = \frac{1}{2}$	A1	
	(iii)	Sub. $x = k \cosh u$	M1	
		Replace all x to $\int k_1 \sinh^2 u \, du$	A1	
		Replace as $\int k_2 (\cosh 2u - 1) \, du$	M1	Or exponential equivalent
		Integrate correctly	A1√	No need for c
		Attempt to replace u with x equivalent	M1	In their answer
		Tidy to reasonable form	A1	cao ($\frac{1}{2}x\sqrt{4x^2 - 1} - \frac{1}{4} \cosh^{-1} 2x (+c)$)