



**ADVANCED GCE UNIT
MATHEMATICS**

Core Mathematics 4

THURSDAY 14 JUNE 2007

4724/01

Afternoon

Time: 1 hour 30 minutes

Additional Materials: Answer Booklet (8 pages)
List of Formulae (MF1)

INSTRUCTIONS TO CANDIDATES

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a graphical calculator in this paper.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is 72.

ADVICE TO CANDIDATES

- Read each question carefully and make sure you know what you have to do before starting your answer.
- **You are reminded of the need for clear presentation in your answers.**

This document consists of **4** printed pages.

2

- 1 The equation of a curve is $y = f(x)$, where $f(x) = \frac{3x + 1}{(x + 2)(x - 3)}$.
- (i) Express $f(x)$ in partial fractions. [2]
- (ii) Hence find $f'(x)$ and deduce that the gradient of the curve is negative at all points on the curve. [3]
- 2 Find the exact value of $\int_0^1 x^2 e^x dx$. [6]
- 3 Find the exact volume generated when the region enclosed between the x -axis and the portion of the curve $y = \sin x$ between $x = 0$ and $x = \pi$ is rotated completely about the x -axis. [6]
- 4 (i) Expand $(2 + x)^{-2}$ in ascending powers of x up to and including the term in x^3 , and state the set of values of x for which the expansion is valid. [5]
- (ii) Hence find the coefficient of x^3 in the expansion of $\frac{1 + x^2}{(2 + x)^2}$. [2]
- 5 A curve C has parametric equations
- $$x = \cos t, \quad y = 3 + 2 \cos 2t, \quad \text{where } 0 \leq t \leq \pi.$$
- (i) Express $\frac{dy}{dx}$ in terms of t and hence show that the gradient at any point on C cannot exceed 8. [4]
- (ii) Show that all points on C satisfy the cartesian equation $y = 4x^2 + 1$. [3]
- (iii) Sketch the curve $y = 4x^2 + 1$ and indicate on your sketch the part which represents C . [2]
- 6 The equation of a curve is $x^2 + 3xy + 4y^2 = 58$. Find the equation of the normal at the point $(2, 3)$ on the curve, giving your answer in the form $ax + by + c = 0$, where a , b and c are integers. [8]
- 7 (i) Find the quotient and the remainder when $2x^3 + 3x^2 + 9x + 12$ is divided by $x^2 + 4$. [4]
- (ii) Hence express $\frac{2x^3 + 3x^2 + 9x + 12}{x^2 + 4}$ in the form $Ax + B + \frac{Cx + D}{x^2 + 4}$, where the values of the constants A , B , C and D are to be stated. [1]
- (iii) Use the result of part (ii) to find the exact value of $\int_1^3 \frac{2x^3 + 3x^2 + 9x + 12}{x^2 + 4} dx$. [5]

3

- 8 The height, h metres, of a shrub t years after planting is given by the differential equation

$$\frac{dh}{dt} = \frac{6-h}{20}.$$

A shrub is planted when its height is 1 m.

- (i) Show by integration that $t = 20 \ln\left(\frac{5}{6-h}\right)$. [6]
- (ii) How long after planting will the shrub reach a height of 2 m? [1]
- (iii) Find the height of the shrub 10 years after planting. [2]
- (iv) State the maximum possible height of the shrub. [1]
- 9 Lines L_1 , L_2 and L_3 have vector equations

$$L_1: \mathbf{r} = (5\mathbf{i} - \mathbf{j} - 2\mathbf{k}) + s(-6\mathbf{i} + 8\mathbf{j} - 2\mathbf{k}),$$

$$L_2: \mathbf{r} = (3\mathbf{i} - 8\mathbf{j}) + t(\mathbf{i} + 3\mathbf{j} + 2\mathbf{k}),$$

$$L_3: \mathbf{r} = (2\mathbf{i} + \mathbf{j} + 3\mathbf{k}) + u(3\mathbf{i} + c\mathbf{j} + \mathbf{k}).$$

- (i) Calculate the acute angle between L_1 and L_2 . [4]
- (ii) Given that L_1 and L_3 are parallel, find the value of c . [2]
- (iii) Given instead that L_2 and L_3 intersect, find the value of c . [5]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (OCR) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

OCR is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.